

This article appeared in a journal published by Elsevier. The attached copy is furnished to the author for internal non-commercial research and education use, including for instruction at the authors institution and sharing with colleagues.

Other uses, including reproduction and distribution, or selling or licensing copies, or posting to personal, institutional or third party websites are prohibited.

In most cases authors are permitted to post their version of the article (e.g. in Word or Tex form) to their personal website or institutional repository. Authors requiring further information regarding Elsevier's archiving and manuscript policies are encouraged to visit:

<http://www.elsevier.com/copyright>



Contents lists available at SciVerse ScienceDirect

## Statistics and Probability Letters

journal homepage: [www.elsevier.com/locate/stapro](http://www.elsevier.com/locate/stapro)

## A general measure of skewness

Magnus Ekström<sup>a</sup>, Sreenivasa Rao Jammalamadaka<sup>b,\*</sup><sup>a</sup> Centre of Biostochastics, Swedish University of Agricultural Sciences, Umeå, Sweden<sup>b</sup> Department of Statistics and Applied Probability, University of California, Santa Barbara, USA

## ARTICLE INFO

## Article history:

Received 9 December 2011

Received in revised form 11 April 2012

Accepted 11 April 2012

Available online 21 April 2012

## Keywords:

Measure of skewness

Bowley's coefficient of skewness

Test of symmetry

Large sample theory

Asymptotic relative efficiency

## ABSTRACT

A very general measure of skewness based on the quantiles is introduced, which includes several well-known measures as special cases. Sample versions of our measure may be used as test statistics for testing the hypothesis of symmetry about an unknown value. We provide large sample theory for such a statistic and discuss the asymptotic relative efficiencies of this against some competing test statistics for symmetry.

© 2012 Elsevier B.V. All rights reserved.

## 1. Introduction

For symmetric unimodal distributions, the three most common measures of central tendency, the mean  $\mu$ , the median  $M$ , and the mode  $m$ , coincide. For other distributions, the lack of symmetry is expressed through measures of skewness. The idea of measuring skewness of a distribution has roots going back at least to Pearson (1895), who suggested the measure  $(\mu - m)/\sigma$ , where  $\sigma$  is the standard deviation of the distribution. Pearson (1895) also used a quantity denoted by  $\beta_1$ , which is the square of the standardized third moment, and Charlier (1906) proposed  $-6^{-1}\beta_1^{1/2}$  as a measure of skewness. Today, the standardized third moment

$$\gamma_1 = \beta_1^{1/2} = \mu_3/\sigma^3,$$

where  $\mu_k$  is the  $k$ th central moment which is often referred to as “the coefficient of skewness”, although any odd-moment can be used for the purpose. Other early measures include Bowley's (1901) coefficient of skewness,

$$S_{\text{Bowley}} = \frac{Q_{0.75} - 2M + Q_{0.25}}{Q_{0.75} - Q_{0.25}},$$

where  $Q_{0.25}$  and  $Q_{0.75}$  represent the 25th and 75th percentiles, and the measure

$$S_{\text{Yule}} = \frac{\mu - M}{\sigma},$$

given in Yule (1911).

A generalization of Bowley's measure of skewness was presented by David and Johnson (1956),

$$S_{\text{D\&J}} = \frac{Q_{1-\alpha} - 2M + Q_{\alpha}}{Q_{1-\alpha} - Q_{\alpha}},$$

\* Corresponding author. Tel.: +1 805 8933119; fax: +1 805 8932334.

E-mail addresses: [Magnus.Ekstrom@slu.se](mailto:Magnus.Ekstrom@slu.se) (M. Ekström), [rao@pstat.ucsb.edu](mailto:rao@pstat.ucsb.edu) (S.R. Jammalamadaka).

for any  $\alpha$  between 0 and  $1/2$  (see also [Hinkley, 1975](#)). Note that Bowley's measure (or coefficient) of skewness is obtained by setting  $\alpha = 0.25$ . The measure obtained by setting  $\alpha = 0.1$  is often referred to as Kelly's coefficient of skewness. By integrating out  $\alpha$ , [Groeneveld and Meeden \(1984\)](#) obtained another measure of skewness viz.

$$S_{G\&M} = \frac{\int_0^{1/2} (Q_{1-\alpha} - 2M + Q_\alpha) d\alpha}{\int_0^{1/2} (Q_{1-\alpha} - Q_\alpha) d\alpha} = \frac{\mu - M}{E|X - M|}$$

sample version of which may be regarded as a robustified version of the sample analogue of  $S_{Yule}$  ([Miao et al., 2006](#)). Bowley's as well as the skewness measures  $S_{Yule}$ ,  $S_{D\&J}$ ,  $S_{G\&M}$ , are bounded by 1 on their absolute values (see, e.g., [Groeneveld, 1991](#)), which may be considered an advantage.

In the current paper we introduce a very general measure of skewness based on the quantiles, that contains the measures by Bowley, Kelly, and David and Johnson as special cases. Also a particular sample version of our measure, when using as many quantiles as the sample size, leads to the sample version of Groeneveld and Meeden's measure, and a test statistic proposed by [Miao et al. \(2006\)](#).

Sample versions of our measure may be used as test statistics for testing the hypothesis of symmetry about an unknown value. We provide results for the asymptotic distribution of such test statistics under the null hypothesis of symmetry, as well as under a sequence of converging alternatives of the form  $F(t + n^{-1/2}\gamma(t))$ , where the distribution function  $F$  is symmetric about zero,  $\gamma$  is some smooth function, and  $n$  is the sample size. This allows us to make comparisons of the Asymptotic Relative Efficiencies (AREs) of these competing tests, which we do in Section 3. A numerical example from forestry is given in Section 4.

## 2. Main results

Suppose  $X_1, \dots, X_n$  are independently and identically distributed (i.i.d.) real-valued random variables with cumulative distribution function (cdf)  $F$ . For any  $0 < \alpha < 1$ , the population quantile function is given by

$$F^{-1}(\alpha) = \inf\{x : F(x) \geq \alpha\},$$

and will also be denoted as  $Q_\alpha$ . The empirical cdf is denoted as  $F_n$ , while

$$F_n^{-1}(\alpha) = \inf\{x : F_n(x) \geq \alpha\}$$

will denote the empirical quantile function, for which we will also use the notation  $\hat{Q}_\alpha = F_n^{-1}(\alpha)$ .

We now quote a result on the so-called Bahadur representation of quantiles:

**Lemma 1** ([Ghosh, 1971](#)). Let  $0 < \alpha < 1$ . Suppose  $F$  is differentiable at  $Q_\alpha$  with  $F'(Q_\alpha) > 0$ . Then

$$\hat{Q}_\alpha - Q_\alpha = \frac{\alpha - F_n(Q_\alpha)}{F'(Q_\alpha)} + o_p(n^{-1/2}). \quad (1)$$

For an integer  $r \geq 2$ , let  $\alpha_1, \dots, \alpha_{2r-1}$  be a finite sequence of real numbers such that

$$0 < \alpha_1 < \alpha_2 < \dots < \alpha_r = 1/2 < \alpha_{r+1} < \dots < \alpha_{2r-1} < 1$$

and  $1 - \alpha_i = \alpha_{2r-i}$ ,  $i = 1, \dots, r$ . Set  $f_j = F'(Q_{\alpha_j})$ ,  $j = 1, \dots, 2r - 1$ . The representation in (1) and the multivariate central limit theorem imply that  $n^{1/2}(\hat{Q}_{\alpha_1} - Q_{\alpha_1}, \dots, \hat{Q}_{\alpha_{2r-1}} - Q_{\alpha_{2r-1}})$  is asymptotically normal with a zero mean vector and covariances  $\text{cov}_{ij} = \alpha_i(1 - \alpha_j)/(f_i f_j)$  for  $i \leq j$ , and  $\text{cov}_{ij} = \text{cov}_{ji}$  for  $i > j$ .

We now define our general measure of skewness,

$$S_r = \frac{\sum_{j=1}^{r-1} ((Q_{\alpha_{2r-j}} - Q_{\alpha_r}) - (Q_{\alpha_r} - Q_{\alpha_j}))}{\sum_{j=1}^{r-1} ((Q_{\alpha_{2r-j}} - Q_{\alpha_r}) + (Q_{\alpha_r} - Q_{\alpha_j}))} = \frac{\sum_{j=1}^{2r-1} c_j Q_{\alpha_j}}{\sum_{j=1}^{2r-1} d_j Q_{\alpha_j}},$$

where  $c_i = -2(r - 1)$  if  $i = r$  and 1 otherwise, and  $d_i = -1$  if  $i < r$ , 0 if  $i = r$ , and 1 if  $i > r$ . We will call the numerator and denominator  $A_r = \sum_{j=1}^{2r-1} c_j Q_{\alpha_j}$ , and  $B_r = \sum_{j=1}^{2r-1} d_j Q_{\alpha_j}$  respectively, and denote the sample analogue of  $S_r$  by  $\hat{S}_{r,n}$ .

**Theorem 1.** Suppose  $F$  is differentiable at  $Q_{\alpha_j}$ ,  $j = 1, \dots, 2r - 1$ . If  $f_j > 0$  for all  $j$ , then

$$\sqrt{n}(\hat{S}_{r,n} - S_r) \xrightarrow{d} N(0, \tau^2),$$

where

$$\tau^2 = \frac{B_r^2 \tau_1^2 - 2A_r B_r \tau_{12} + A_r^2 \tau_2^2}{B_r^4}$$

and

$$\begin{aligned}\tau_1^2 &= \sum_{i=1}^{2r-1} \sum_{j=1}^{2r-1} \frac{c_i c_j \alpha_{\min\{i,j\}} (1 - \alpha_{\max\{i,j\}})}{f_i f_j}, \\ \tau_{12} &= \sum_{i=1}^{2r-1} \sum_{j=1}^{2r-1} \frac{c_i d_j \alpha_{\min\{i,j\}} (1 - \alpha_{\max\{i,j\}})}{f_i f_j}, \\ \tau_2^2 &= \sum_{i=1}^{2r-1} \sum_{j=1}^{2r-1} \frac{d_i d_j \alpha_{\min\{i,j\}} (1 - \alpha_{\max\{i,j\}})}{f_i f_j}.\end{aligned}$$

**Proof.** Let  $\hat{A}_{r,n}$  and  $\hat{B}_{r,n}$  be the sample analogues of  $A_r$  and  $B_r$ , respectively. Then

$$\sqrt{n}(\hat{S}_{r,n} - S_r) = \sqrt{n} \cdot \frac{B_r(\hat{A}_{r,n} - A_r) - A_r(\hat{B}_{r,n} - B_r)}{\hat{B}_{r,n} B_r}. \quad (2)$$

By Lemma 1,

$$\hat{A}_{r,n} - A_r = \bar{Y} + o_p(n^{-1/2}),$$

where  $\bar{Y}$  is the mean value of

$$Y_i = \sum_{j=1}^{2r-1} \frac{c_j(\alpha_j - I(X_i \leq Q_{\alpha_j}))}{f_j}, \quad i = 1, \dots, n, \quad (3)$$

and

$$\hat{B}_{r,n} - B_r = \bar{Z} + o_p(n^{-1/2}),$$

where  $\bar{Z}$  is the mean value of

$$Z_i = \sum_{j=1}^{2r-1} \frac{d_j(\alpha_j - I(X_i \leq Q_{\alpha_j}))}{f_j}, \quad i = 1, \dots, n.$$

By the multivariate central limit theorem,  $n^{1/2}\bar{Y}$  and  $n^{1/2}\bar{Z}$  are jointly asymptotically normal with mean zero and covariance matrix

$$\begin{pmatrix} \tau_1^2 & \tau_{12} \\ \tau_{12} & \tau_2^2 \end{pmatrix}.$$

By Slutsky's theorem, the same is true for  $n^{1/2}(\hat{A}_{r,n} - A_r)$  and  $n^{1/2}(\hat{B}_{r,n} - B_r)$ , which implies that

$$\sqrt{n}(B_r(\hat{A}_{r,n} - A_r) - A_r(\hat{B}_{r,n} - B_r)) \xrightarrow{d} N(0, \tau^2). \quad (4)$$

The conclusion of the theorem follows from (2), (4), and the Slutsky's theorem by noting that  $\hat{B}_{r,n}$  tends in probability to  $B_r$  as  $n \rightarrow \infty$ .  $\square$

We consider now the null hypothesis of symmetry about an unknown point of symmetry. Under the assumption that the distribution function  $F$  is symmetric about some unknown value  $\mu$ , i.e.  $F(t - \mu) = 1 - F(-(t - \mu))$  for all  $t$ , we see that Theorem 1 simplifies to

$$\sqrt{n}(\hat{S}_{r,n} - S_r) \xrightarrow{d} N(0, B_r^{-2} \tau_1^2).$$

The asymptotic distribution of  $\hat{S}_{r,n}$  under close alternatives is studied next. As in Antille and Kersting (1977) and Ekström and Jammalamadaka (2007), we consider close alternatives of the form  $G_n(t) = F(t + n^{-1/2}\gamma(t))$ . The function  $t + n^{-1/2}\gamma(t)$  is assumed to be a monotonically increasing function of  $t$  for large  $n$ . We remark that a result of Wang (2008) deals with the special case where  $r = 2$  and  $\gamma(t) = \gamma_n(t) = -\eta t / (1 + n^{-1/2}\eta)$  if  $t > 0$ , and 0 otherwise.

**Theorem 2.** Assume, without loss of generality, that  $F$  is symmetric about 0, that the density function  $f$  is twice continuously differentiable and strictly positive on the whole real line, and that the function  $\gamma$  is continuously differentiable. Let

$$W_i = \frac{\gamma(X_i)f'(X_i)}{f(X_i)} + \gamma'(X_i),$$

and assume that  $E_{H_0} W_1^2$  and  $E_{H_0} \gamma^2(X_1)$  are finite, and that  $\lim_{t \rightarrow \pm\infty} \gamma(t)f(t) = \lim_{t \rightarrow \pm\infty} \gamma^2(t)f'(t) = 0$ . Then, under  $H_{A,n}$ ,

$$n^{1/2} \hat{S}_{r,n} \xrightarrow{d} N(B_r^{-1} E_{H_0} Y_1 W_1, B_r^{-2} \tau_1^2),$$

where  $Y_1$  is defined in (3).

**Proof.** By Le Cam's third lemma (see, e.g., van der Vaart, 1998), we can obtain the limiting distribution of  $n^{1/2} \hat{S}_{r,n}$  under  $H_{A,n}$ , once we have the joint limit distribution of  $L_n = \sum_{i=1}^n \log(G'_n(X_i)/F'(X_i))$  and  $n^{1/2} \hat{S}_{r,n}$  under  $H_0$ . We begin with  $L_n$ . Let

$$V_i = \frac{f(X_i + n^{-1/2} \gamma(X_i)) - f(X_i) + n^{-1/2} \gamma'(X_i) f(X_i + n^{-1/2} \gamma(X_i))}{f(X_i)}.$$

If we write  $\log(1+x) = x - x^2/2 + x^2 R(x)$ , then  $R(x) \rightarrow 0$  as  $x \rightarrow 0$ , and

$$\begin{aligned} L_n &= \sum_{i=1}^n \log \left( \frac{(1 + n^{-1/2} \gamma'(X_i)) f(X_i + n^{-1/2} \gamma(X_i))}{f(X_i)} \right) \\ &= \sum_{i=1}^n \log(1 + V_i) \\ &= \sum_{i=1}^n V_i - \frac{1}{2} \sum_{i=1}^n V_i^2 + \sum_{i=1}^n V_i^2 R(V_i). \end{aligned}$$

By Taylor expansions,

$$\begin{aligned} \sum_{i=1}^n V_i &= \frac{1}{n^{1/2}} \sum_{i=1}^n \frac{\gamma(X_i) f'(X_i)}{f(X_i)} + \frac{1}{2n} \sum_{i=1}^n \frac{\gamma^2(X_i) f''(X_i)}{f(X_i)} + \frac{1}{2n} \sum_{i=1}^n \frac{\gamma^2(X_i) (f''(X_i + n^{-1/2} \rho_i \gamma(X_i)) - f''(X_i))}{f(X_i)} \\ &\quad + \frac{1}{n^{1/2}} \sum_{i=1}^n \gamma'(X_i) + \frac{1}{n} \sum_{i=1}^n \frac{\gamma(X_i) \gamma'(X_i) f'(X_i)}{f(X_i)} \\ &\quad + \frac{1}{n} \sum_{i=1}^n \frac{\gamma(X_i) \gamma'(X_i) (f'(X_i + n^{-1/2} \eta_i \gamma(X_i)) - f'(X_i))}{f(X_i)}, \end{aligned} \quad (5)$$

where all  $\rho_i$  and  $\eta_i$  are between 0 and 1. Consider the third term on the right. For each  $\delta_* > 0$  there exist a closed interval  $[a_1, a_2]$  and an integer  $N > 0$  such that  $P_{H_0}(X_1 \in B_{r,n}) \geq 1 - \delta_*/2$  for all  $n > N$ , where  $B_{r,n} = \{t : a_1 \leq t \leq a_2 \text{ and } a_1 \leq t + n^{-1/2} \gamma(t) \leq a_2\}$ . By our assumptions,  $f''$  is uniformly continuous in  $[a_1, a_2]$ . Hence, for each  $\varepsilon > 0$  there exists a  $\delta > 0$  such that for all  $t$  and  $t + n^{-1/2} \rho_i \gamma(t)$  in  $[a_1, a_2]$  with  $|n^{-1/2} \rho_i \gamma(t)| \leq \delta$ , we have  $|f''(t + n^{-1/2} \rho_i \gamma(t)) - f''(t)| \leq \varepsilon$ . Thus, for all  $n$  large enough,

$$\begin{aligned} &P_{H_0} \left( \max_{1 \leq i \leq n} |(f''(X_i + n^{-1/2} \rho_i \gamma(X_i)) - f''(X_i))| \geq \varepsilon \right) \\ &\leq P_{H_0} \left( \max_{1 \leq i \leq n} |(f''(X_i + n^{-1/2} \rho_i \gamma(X_i)) - f''(X_i))| \geq \varepsilon \text{ and } X_i \in B_{r,n} \right) + P_{H_0}(X_1 \notin B_{r,n}) \\ &\leq P_{H_0} \left( \max_{1 \leq i \leq n} |\gamma(X_i)| \geq n^{1/2} \delta \text{ and } X_i \in B_{r,n} \right) + P_{H_0}(X_1 \notin B_{r,n}) \leq \delta_*, \end{aligned} \quad (6)$$

where the last inequality follows from the fact that  $\gamma$  is a bounded function on  $\bigcup B_{r,n}$ . The third term on the right-hand side of (5) is bounded by

$$\max_{1 \leq i \leq n} |(f''(X_i + n^{-1/2} \rho_i \gamma(X_i)) - f''(X_i))| \frac{1}{2n} \sum_{j=1}^n \frac{\gamma^2(X_j)}{f(X_j)}.$$

Thus it is  $o_p(1)O_p(1)$ , and converges in probability to zero. Similarly, the last term on the right-hand side of (5) tends to zero in probability. Thus, under  $H_0$  we have

$$\sum_{i=1}^n V_i = \frac{1}{n^{1/2}} \sum_{i=1}^n \frac{\gamma(X_i) f'(X_i)}{f(X_i)} + \frac{1}{n^{1/2}} \sum_{i=1}^n \gamma'(X_i) + \frac{1}{2} E_{H_0} \frac{\gamma^2(X_1) f''(X_1)}{f(X_1)} + E_{H_0} \frac{\gamma(X_1) \gamma'(X_1) f'(X_1)}{f(X_1)} + o_p(1),$$

and, by similar arguments as above,

$$\sum_{i=1}^n V_i^2 = E_{H_0} \left( \frac{\gamma(X_1) f'(X_1)}{f(X_1)} \right)^2 + E_{H_0} (\gamma'(X_1))^2 + 2E_{H_0} \frac{\gamma(X_1) \gamma'(X_1) f'(X_1)}{f(X_1)} + o_p(1).$$

By arguments similar to those in (6), we see that  $\max_{1 \leq i \leq n} |V_i|$  converges in probability to zero. By the definition of the function  $R$ , the sequence  $\max_{1 \leq i \leq n} |R(V_i)|$  converges in probability to zero as well. Combining these results, we obtain that

$$\begin{aligned} L_n &= \frac{1}{n^{1/2}} \sum_{i=1}^n \frac{\gamma(X_i) f'(X_i)}{f(X_i)} + \frac{1}{n^{1/2}} \sum_{i=1}^n \gamma'(X_i) + \frac{1}{2} E_{H_0} \frac{\gamma^2(X_1) f''(X_1)}{f(X_1)} \\ &\quad - \frac{1}{2} E_{H_0} \left( \frac{\gamma(X_1) f'(X_1)}{f(X_1)} \right)^2 - \frac{1}{2} E_{H_0} (\gamma'(X_1))^2 + o_p(1) \\ &= \frac{1}{n^{1/2}} \sum_{i=1}^n W_i - \frac{1}{2} E_{H_0} W_1^2 + o_p(1), \end{aligned} \quad (7)$$

where the last equality follows by integration by parts. By another integration by parts we see that  $E_{H_0} W_1 = 0$ .

By the proof of Theorem 1,  $n^{1/2} \hat{S} = n^{1/2} B_r^{-1} \bar{Y} + o_p(1)$  under  $H_0$ . Thus, under  $H_0$ ,  $n^{1/2} \hat{S}_{r,n}$  and  $L_n$  are jointly asymptotically normal,

$$N \left( \begin{pmatrix} 0 \\ -2^{-1} E_{H_0} W_1^2 \end{pmatrix}, \begin{pmatrix} B_r^{-2} \tau_1^2 & B_r^{-1} E_{H_0} Y_1 W_1 \\ B_r^{-1} E_{H_0} Y_1 W_1 & E_{H_0} W_1^2 \end{pmatrix} \right),$$

and under  $H_{A,n}$ , Le Cam's third lemma implies that

$$n^{1/2} \hat{S}_{r,n} \xrightarrow{d} N(B_r^{-1} E_{H_0} Y_1 W_1, B_r^{-2} \tau_1^2). \quad \square$$

In Theorems 1 and 2, a fixed number,  $2r - 1$ , of sample quantiles have been used to define the statistic  $\hat{S}_{r,n}$ . Next we consider a case where this number is allowed to increase with the sample size. That is, we will consider the case where  $2r - 1 = n$ , so that we have as many sample quantiles defining the statistic  $\hat{S}_{r,n}$  as observations in the sample. Assume, for simplicity, that  $n = 2r - 1$  is an odd number, and set  $\alpha_i = i/(n + 1)$ ,  $i = 1, \dots, n$ . Then  $\hat{Q}_{\alpha_i} = X_{(i)}$  for each  $i$ , where  $X_{(i)}$  is the  $i$ th order statistic of the sample, and our statistic can be written in the following form,

$$\hat{S}_{r,n} = \frac{\bar{X} - \hat{M}}{\hat{D}_n}, \quad (8)$$

where  $r_n = (n + 1)/2$ ,  $\hat{M}$  is the sample median, and  $\hat{D}_n = n^{-1} \sum_{j=1}^n |X_j - \hat{M}|$ .

Let  $\mu$ ,  $M$  and  $\sigma^2$  denote respectively the mean, median and variance of the distribution  $F$ . Let  $D = E|X_1 - M|$ ,

$$v^2 = \sigma^2 + \frac{1}{4f^2(M)} - \frac{D}{f(M)},$$

and

$$U_i = (X_i - \mu) - \frac{1/2 - I(X_i \leq M)}{f(M)}, \quad i = 1, \dots, n. \quad (9)$$

**Theorem 3.** Assume that  $\sigma^2 < \infty$  and that the assumptions of Theorem 2 are valid. Then, under  $H_{A,n}$ ,

$$n^{1/2} \hat{S}_{r,n} \xrightarrow{d} N(D^{-1} E_{H_0} U_1 W_1, D^{-2} v^2),$$

where  $\hat{S}_{r,n}$  is defined as in (8).

**Proof.** Again, with the help of Le Cam's third lemma, we will obtain the limit distribution of  $n^{1/2} \hat{S}_{r,n}$  under  $H_{A,n}$ , once the joint limit distribution of  $L_n = \sum_{i=1}^n \log(G'_n(X_i)/F'(X_i))$  and  $n^{1/2} \hat{S}_{r,n}$  under  $H_0$  is known.

By Lemma 1, under  $H_0$ ,

$$\bar{X} - \hat{M} = (\bar{X} - \mu) - (\hat{M} - M) = \bar{U} + o_p(n^{-1/2}),$$

where  $\bar{U}$  is the mean value of  $U_1, \dots, U_n$ . By the central limit theorem,  $n^{1/2} \bar{U}$  is, under  $H_0$ , asymptotically normal with mean zero and variance  $v^2$ , and by Slutsky's theorem, the same is true for  $n^{1/2} (\bar{X} - \hat{M})$ . The statistic  $\hat{D}_n$  converges in probability to  $D$  as  $n \rightarrow \infty$  (Mira, 1999). Thus, under  $H_0$ ,  $n^{1/2} \hat{S}_{r,n} = n^{1/2} D^{-1} \bar{U} + o_p(1)$ . Under  $H_0$ , this result combined with (7) imply that  $n^{1/2} \hat{S}_{r,n}$  and  $L_n$  are jointly asymptotically normal,

$$N \left( \begin{pmatrix} 0 \\ -2^{-1} E_{H_0} W_1^2 \end{pmatrix}, \begin{pmatrix} D^{-2} v^2 & D^{-1} E_{H_0} U_1 W_1 \\ D^{-1} E_{H_0} U_1 W_1 & E_{H_0} W_1^2 \end{pmatrix} \right),$$

and under  $H_{A,n}$ , Le Cam's third lemma implies that

$$n^{1/2}\hat{S}_{r,n} \xrightarrow{d} N(D^{-1}E_{H_0}U_1W_1, D^{-2}v^2). \quad \square$$

**Remark 1.** The statistic (8),  $\hat{S}_{r,n}$ , is the sample analogue of Groeneveld and Meeden's (1984) measure of skewness, and is essentially the test statistic proposed by Miao et al. (2006), i.e., their statistic is obtained by multiplying the right-hand side of (8) by  $(2/\pi)^{1/2}$ . Miao et al. show, under general conditions, that

$$n^{1/2}\hat{T}_n = n^{1/2} \left( \frac{\bar{X} - \hat{M}}{\hat{D}_n} - \frac{\mu - M}{D} \right)$$

tends to a normal distribution with mean zero and variance

$$\frac{1}{D^2} \left( \sigma^2 + \frac{1}{4f^2(M)} - \frac{D}{f(M)} + \frac{(\mu - M)^2(\sigma^2 + (\mu - M)^2)}{D^2} - (\mu - M)^2 \right. \\ \left. - \frac{2(\mu - M)}{D} (\sigma^2 - 2EX_1^2I(X_1 \leq M) + 2\mu EX_1I(X_1 \leq M) - MD) + \frac{(\mu - M)^2}{Df(M)} \right).$$

Under the null hypothesis when  $F$  is symmetric, then the asymptotic distribution of  $n^{1/2}\hat{T}_n$  is normal with mean zero and variance given by  $D^{-2}v^2$ .

**Remark 2.** Consider the measure of skewness defined as  $S_{Yule} = (\mu - M)/\sigma$ . The sample version of this measure,  $\hat{S}_{Yule,n} = (\bar{X} - \hat{M})/s$ , where  $s$  is the sample standard deviation, is discussed in Gastwirth (1971), and in the case  $\mu = M$ , Cabilio and Masaro (1996) showed that  $n^{1/2}\hat{S}_{Yule,n}$  is asymptotically normal with mean zero and variance  $\sigma^{-2}v^2$ . Under  $H_{A,n}$ , and the assumptions of Theorem 3, it is easily seen that

$$n^{1/2}\hat{S}_{Yule,n} \xrightarrow{d} N(\sigma^{-1}E_{H_0}U_1W_1, \sigma^{-2}v^2).$$

**Remark 3.** Consider the standardized third moment  $\gamma_1 = \mu_3/\sigma^3$ , where  $\mu_k$  is the  $k$ th central moment. The sample version of this measure is  $\hat{\gamma}_{1,n} = m_3/s^3$ , where  $m_k$  is the sample  $k$ th central moment. If  $F$  is symmetric and  $\mu_6 < \infty$ , then  $\gamma_1 = 0$  is zero and  $n^{1/2}\hat{\gamma}_{1,n}$  is asymptotically normal with mean zero and variance  $\sigma^{-6}(\mu_6 - 6\sigma^2\mu_4 + 9\sigma^6)$  (see, e.g., Gupta (1967) for more details). If  $\mu_6 < \infty$  and the assumptions of Theorem 2 are valid, then it can be shown under  $H_{A,n}$  that

$$n^{1/2}\hat{\gamma}_{1,n} \xrightarrow{d} N(\sigma^{-3}(E_{H_0}X_1^3W_1 - 3E_{H_0}X_1^2E_{H_0}X_1W_1), \sigma^{-6}(\mu_6 - 6\sigma^2\mu_4 + 9\sigma^6)).$$

### 3. Pitman asymptotic relative efficiency

The Pitman asymptotic relative efficiency (ARE) of a test relative to another test is defined to be the limit of the inverse ratio of sample sizes required to obtain the same limiting power at a sequence of alternatives converging to the hypothesis. The limiting power should be a value between the limiting test size,  $\alpha$ , and the maximum power, 1. If the limiting power of a test at a sequence of alternatives is  $\alpha$ , then its ARE with respect to any other test with the same test size and with limiting power greater than  $\alpha$ , is zero. On the other hand, if the limiting power of a test at a sequence of alternatives converges to a number in the open interval  $(\alpha, 1)$ , then a measure of rate of convergence, called efficacy, can be computed. Under certain standard regularity assumptions (see, for example, Rao, 1973, p. 469), which include a condition about the nature of the alternative, asymptotic normal distribution of the test statistic under the sequence of alternatives, etc., this efficacy is given by

$$\text{efficacy} = \frac{\mu_*^2}{\sigma_*^2}.$$

Here  $\mu_*$  and  $\sigma_*^2$  are the mean and variance of the limiting normal distribution under the sequence of alternatives when the test statistic has been normalized to have a limiting standard normal distribution under the hypothesis. In such a situation, the ARE of one test with respect to another is simply the ratio of their efficacies.

The efficacy of the test statistic  $n^{1/2}\hat{S}_{r,n}$  from Theorem 2 is

$$\frac{(E_{H_0}Y_1W_1)^2}{\tau_1^2},$$

and the efficacy of the test statistic  $n^{1/2}\hat{S}_{r,n}$  from Theorem 3 is

$$\frac{(E_{H_0}U_1W_1)^2}{v^2}.$$



**Table 1**  
Best choice of quantiles for various distributions.

Distribution	$\theta$	$r$	$\alpha_1, \dots, \alpha_{r-1}$	Max efficacy
$t_1$ (Cauchy)	1	2	0.250	0.101
$t_1$ (Cauchy)	1	3	0.215, 0.329	0.113
$t_1$ (Cauchy)	1	4	0.196, 0.290, 0.361	0.118
$t_2$	2	2	0.167	0.150
$t_2$	2	3	0.128, 0.251	0.170
$t_2$	2	4	0.109, 0.204, 0.288	0.178
EP (Laplace)	1	2	0.102	0.162
EP (Laplace)	1	3	0.067, 0.200	0.195
EP (Laplace)	1	4	0.050, 0.150, 0.250	0.210
EP (normal)	2	2	0.053	0.250
EP (normal)	2	3	0.026, 0.108	0.292
EP (normal)	2	4	0.017, 0.065, 0.140	0.308
Logistic		2	0.085	0.212
Logistic		3	0.051, 0.158	0.248
Logistic		4	0.036, 0.108, 0.193	0.261

Also, it should be noted that the test statistic  $n^{1/2}\hat{S}_{\text{vule},n}$  has the same efficacy as  $n^{1/2}\hat{S}_{r,n}$ . The efficacy of  $n^{1/2}\hat{\gamma}_{1,n}$  from Remark 3 is

$$\frac{(E_{H_0}X_1^3W_1 - 3E_{H_0}X_1^2E_{H_0}X_1W_1)^2}{\mu_6 - 6\sigma^2\mu_4 + 9\sigma^6}.$$

Computations of efficacy will be made in the case where  $\gamma(t) = t$ ,  $t \geq 0$ , and 0 elsewhere. In this case,  $B_r^{-1}E_{H_0}Y_1W_1 = D^{-1}E_{H_0}U_1W_1 = -1/2$ ,  $E_{H_0}X_1^3W_1 = -(3/2)E_{H_0}|X_1|^3$ , and  $E_{H_0}X_1W_1 = -(1/2)E_{H_0}|X_1|$ .

We will consider distributions  $F$  with the following densities,

$$t \text{ distribution : } f_\theta(x) = \frac{\Gamma(\frac{\theta+1}{2})}{(\theta\pi)^{1/2}\Gamma(\frac{\theta}{2})} \left(1 + \frac{x^2}{\theta}\right)^{-\frac{\theta+1}{2}},$$

$$\text{Exponential power (EP) distribution : } f_\theta(x) = \frac{e^{-|x|^\theta}}{2\Gamma(1 + \theta^{-1})},$$

$$\text{Logistic distribution : } f(x) = \frac{e^{-x}}{(1 + e^{-x})^2}.$$

The double exponential distribution, also known as the Laplace distribution, is obtained by setting  $\theta = 1$  in the EP distribution, and the normal distribution with mean zero and variance  $1/2$  is obtained by setting  $\theta = 2$  in the EP distribution. The  $t$  distribution with  $\theta = 1$  degree of freedom is also known as the standard Cauchy distribution.

In Table 1, for each given  $r$ , we choose the  $\alpha_i$ 's which maximize the efficacy. From this table as well as heuristically, it appears that the efficacy increases when more quantiles are brought into play, but a reasonably practical approach may be to use  $r = 3$  with 2 quantiles from either tail. It can also be observed that in thin-tailed distributions like the normal, one should use much smaller-order quantiles e.g. the 5th and 95th percentiles with  $r = 2$ , whereas in fat-tailed models like the Cauchy, Bowley's choice of the 25th and 75th seems optimal. Thus Bowley's is a better choice than Kelly's for heavy tailed distributions whereas the reverse is true for thin-tailed distributions like the normal. In general there is no universally optimal choice of these quantiles for all distributions. For instance we have verified that with  $r = 3$ , by choosing  $\alpha_1 = 1/6$  and  $\alpha_2 = 2/6$  we can do uniformly better than Bowley's test and by choosing  $\alpha_1 = 1/10$  and  $\alpha_2 = 2/10$  we can beat Kelly's test for all the distributions checked.

However if one were to build a reasonably robust test of symmetry with  $r = 3$  which does better over a wide range of distributions, an overall compromise seems to be to take  $\alpha_1 = 0.1$  and  $\alpha_2 = 0.2$  i.e. compare the 10th, 20th, 80th, and 90th percentiles to the median, getting the test:

$$\hat{S}_{3,n}^* = \frac{Q_{0.90} + Q_{0.80} - 4M + Q_{0.20} + Q_{0.10}}{(Q_{0.90} - Q_{0.10}) + (Q_{0.80} - Q_{0.20})}.$$

Such a test as Table 2 indicates, will do just as well or better than Kelly's in all the cases and falls short of Bowley's only in the case of one distribution viz. the Cauchy. Even further improvements over the Bowley's and Kelly's coefficients can be demonstrated if we use  $\hat{S}_{r,n}$  with a larger  $r$  but we omit such comparisons.

In Antille and Kersting (1977), the following four test statistics for symmetry were considered,

$$\hat{S}_{A\&K,1,n} = \sum_{1 \leq i \leq n} \left( I(X_i - \bar{X} \leq 0) - \frac{1}{2} \right),$$



**Table 2**

AREs of  $\hat{S}_{3,n}^*$  relative to Bowley's and Kelly's coefficients.

Distribution	$\theta$	Bowley		Kelly	
		Efficacy	ARE	Efficacy	ARE
$t_1$ (Cauchy)	1	0.101	0.70	0.051	1.38
$t_2$	2	0.129	1.27	0.129	1.27
EP (Laplace)	1	0.120	1.56	0.162	1.16
EP (normal)	2	0.136	1.73	0.233	1.01
Logistic		0.136	1.67	0.211	1.08

**Table 3**

Efficacies of the test statistics.

Distribution	$\theta$	Efficacy						
		$\hat{S}_{3,n}^*$	$\hat{S}_{r,n}$	$\hat{\gamma}_{1,n}$	$\hat{S}_{A\&K_1,n}$	$\hat{S}_{A\&K_2,n}$	$\hat{S}_{A\&K_3,n}$	$\hat{S}_{A\&K_4,0,n}$
$t_1$ (Cauchy)	1	0.071					0.123	0.056
$t_2$	2	0.164					0.171	0.096
EP (Laplace)	1	0.188	0.250	0.112	0.250	0.188	0.188	0.125
EP (normal)	2	0.236	0.279	0.239	0.279	0.289	0.201	0.143
Logistic		0.227	0.275	0.238	0.275	0.216	0.196	0.134

$$\hat{S}_{A\&K_2,n} = \sum_{1 \leq i \leq j \leq n} \left( I(X_i + X_j \leq 2\bar{X}) - \frac{1}{2} \right),$$

$$\hat{S}_{A\&K_3,n} = \sum_{1 \leq i \leq j \leq n} \left( I(X_i + X_j \leq 2\hat{M}) - \frac{1}{2} \right),$$

$$\hat{S}_{A\&K_4,\varepsilon,n} = \sum_{1+[ \varepsilon n ] \leq i \leq [ \varepsilon n ]} \left( I(D_i - D_{n-i+2} \leq 0) - \frac{1}{2} \right),$$


where  $D_i = X_{(i)} - X_{(i-1)}$ ,  $i = 1, \dots, n$ . In their paper, asymptotic distributions under symmetry is given, and asymptotic distributions under alternatives of the form  $G_n(t) = F(t + n^{-1/2}\gamma(t))$ , where  $F$  is symmetric about 0, are also given. Based on their results, the efficacy of each test may be computed. In Ekström and Jammalamadaka (2007), a modification of  $\hat{S}_{A\&K_4,\varepsilon,n}$  is suggested, and unlike Antille and Kersting's test, the modified test is asymptotically distribution-free. The efficacy of the modified test coincides with that for  $\hat{S}_{A\&K_4,\varepsilon,n}$ .

**Remark 4.** Under the alternative,  $n^{1/2}\hat{S}_{A\&K_4,\varepsilon,n}$  is asymptotically normal with mean  $\mu_{4,\varepsilon}$  and variance  $\sigma_{4,\varepsilon}^2$ , and  $\mu_{4,\varepsilon}$  and  $\sigma_{4,\varepsilon}^2$  possess limits as  $\varepsilon \rightarrow 0$ ,  $\mu_4$  and  $\sigma_4^2$ , respectively, but it is not known whether  $n^{1/2}\hat{S}_{A\&K_4,0,n}$  is asymptotically normal with mean  $\mu_4$  and variance  $\sigma_4^2$ . However, based on the claim of Antille and Kersting that it is, we compute the efficacy of  $n^{1/2}\hat{S}_{A\&K_4,0,n}$  as  $\mu_4^2/\sigma_4^2$ .

Table 3 provides the efficacies of  $\hat{S}_{3,n}^*$  against Antille and Kersting's four tests,  $\hat{S}_{r,n}$  given in Eq. (8), and the classical coefficient of skewness  $\hat{\gamma}_{1,n}$ . It is interesting to note that the proposed  $\hat{S}_{3,n}^*$ , although based on just 5 quantiles and easier to compute, compares reasonably well with all these other tests which use all the observations/quantiles or gaps between them.

#### 4. A practical example from forestry

We consider two data sets taken from Matérn (1981, Table 2.1), consisting of the diameters at breast height in millimeters from year 1912 and year 1951 for  $n = 115$  Norway spruce trees from the Forest Research Institute of Sweden's sample plot number 238 at Finnerödja. As seen in Fig. 1, the data from 1912 show only a slight skewness, while a more distinct (negative) skewness is visible in the data from 1951. Assume that the data (from 1912 or 1951) are taken from some distribution  $F$ , and that we want to use the test statistic  $\hat{S}_{r,n}$  for testing the hypothesis of symmetry, i.e. that  $F(x - \mu) = 1 - F(-(x - \mu))$  for all  $x$ . If  $F$  is symmetric and  $\hat{\tau}_{1,n}$  is a consistent estimator of  $\tau_1$ , then Theorem 1 implies that  $n^{1/2}\hat{S}_{r,n}\hat{B}_{r,n}\hat{\tau}_{1,n}^{-1}$  is approximately  $N(0, 1)$  for large  $n$ , and approximate  $p$ -values of the test can be computed. In what follows we use Kraft et al.'s (1985) symmetrized kernel density estimator for estimating the density values  $f_j$ ,  $j = 1, \dots, 2r - 1$ , in the definition of  $\tau_1$ . The kernel density estimation is performed using the R function "density" (R Development Core Team, 2011) with a Gaussian kernel. (Another possibility for computing approximate  $p$ -values is to use bootstrap; see e.g. Thomas, 2009 and Zheng and Gastwirth, 2010.) For both data sets, the values of  $\hat{S}_{3,n}^*$  and Bowley's and Kelly's coefficients are given in Table 4. Corresponding  $p$ -values are also given. As expected, all  $p$ -values are far from a significance level of 0.05 for the data



**Fig. 1.** Normal probability plots and histograms of  $n = 115$  diameters at breast height from year 1912 and year 1951.

**Table 4**

Values of three coefficients of skewness, and corresponding  $p$ -values, computed from the two sets of diameter data.

Year	Bowley		Kelly		$\hat{S}_{3,n}^*$	
	Value	$p$ -value	Value	$p$ -value	Value	$p$ -value
1912	0.000	1.00	−0.104	0.34	−0.067	0.52
1951	0.030	0.83	−0.285	0.02	−0.224	0.04

from 1912. For the data from 1951, Bowley's coefficient fails to reject the hypothesis of symmetry, with a corresponding  $p$ -value as large as 0.83, whereas the tests based on Kelly's coefficient and  $\hat{S}_{3,n}^*$  do reject the null hypothesis, which corresponds well with the visual inspection of Fig. 1. The failure of Bowley's coefficient in detecting the obvious skewness in the 1951 data in particular, illustrates the danger of relying on a test based only on one quantile from each tail, like the Bowley's and Kelly's.

## 5. Summary

In the current article we propose a very general class of distribution-free tests of symmetry that encompass most known measures based on quantiles. We provide large sample theory for such statistics and compare their asymptotic relative efficiencies against some competing test statistics for symmetry. In particular we suggest using a simple test  $\hat{S}_{3,n}^*$  which uses 5 quantiles and fares well against all the other known tests of symmetry for a wide variety of distributions. Computer code in R for implementing this test, is available from the authors.

## Acknowledgment

We thank the anonymous referee for a careful reading of this paper as well as for the insightful suggestion that we add a practical example to illustrate the proposed statistics.

## References

- Antille, A., Kersting, G., 1977. Tests for symmetry. *Z. Wahrscheinlichkeitstheor. Verwandte Geb.* 39, 235–255.
- Bowley, L., 1901. *Elements of Statistics*, 6th ed. Staples Press Ltd., London, 1937.
- Cabilio, P., Masaro, J., 1996. A simple test of symmetry about an unknown median. *Canad. J. Statist.* 24, 349–361.

- Charlier, C.V.L., 1906. Researches into the Theory of Probability. In: Kongl. Fysiografiska Sällskapetets Handlingar, vol. 16.
- David, F.N., Johnson, N.L., 1956. Some tests of significance with ordered variables. *J. R. Stat. Soc. Ser. B Stat. Methodol.* 18, 1–31.
- Ekström, M., Jammalamadaka, S.R., 2007. An asymptotically distribution-free test of symmetry. *J. Statist. Plann. Inference* 137, 799–810.
- Gastwirth, J.L., 1971. On the sign test of symmetry. *J. Amer. Statist. Assoc.* 66, 821–823.
- Ghosh, J.K., 1971. A new proof of the Bahadur representation of quantiles and an application. *Ann. Math. Statist.* 42, 1957–1961.
- Groeneveld, R.A., 1991. Sharp inequalities for skewness measures. *Statistician* 40, 387–392.
- Groeneveld, R.A., Meeden, G., 1984. Measuring skewness and kurtosis. *Statistician* 33, 391–399.
- Gupta, M.K., 1967. An asymptotically nonparametric test of symmetry. *Ann. Math. Statist.* 38, 849–866.
- Hinkley, D.V., 1975. On power transformations to symmetry. *Biometrika* 62, 101–111.
- Kraft, C., Lepage, Y., van Eeden, C., 1985. Estimation of a symmetric density function. *Comm. Statist. Theory Methods* 14, 273–288.
- Matérn, B., 1981. Sampling, Part 1, second ed. Swedish University of Agricultural Sciences, Section of Forest Biometry, Umeå, Sweden, (in Swedish).
- Miao, W., Gel, Y.R., Gastwirth, J.L., 2006. A new test of symmetry about an unknown median. In: Hsiung, A., Zhang, C.-H., Ying, Z. (Eds.), *Random Walk, Sequential Analysis and Related Topics—A Festschrift in Honor of Yuan-Shih Chow*. World Scientific, Singapore.
- Mira, A., 1999. Distribution-free test for symmetry based on Bonferroni's measure. *J. Appl. Stat.* 26, 959–971.
- Pearson, K., 1895. Contributions to the mathematical theory of evolution II. Skew variation in homogeneous material. *Philos. Trans. R. Soc. Lond. Ser. A* 186, 343–414.
- Rao, C.R., 1973. *Linear Statistical Inference and its Applications*, second ed. John Wiley, New York.
- R Development Core Team, 2011. R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. ISBN: 3-900051-07-0. URL: <http://www.R-project.org/>.
- Thomas, G.E., 2009. Assessing symmetry using quantiles and *L*-moments. *Comm. Statist. Simulation Comput.* 38, 335–354.
- van der Vaart, A.W., 1998. *Asymptotic Statistics*. Cambridge University Press, Cambridge.
- Wang, J., 2008. Some properties of robust statistics under asymmetric models. Ph.D. Thesis. Rutgers, The State University of New Jersey.
- Yule, G.U., 1911. *Introduction to the Theory of Statistics*, eighth ed. Griffin, London, 1927.
- Zheng, T., Gastwirth, J.L., 2010. On bootstrap tests of symmetry about an unknown median. *J. Data Sci.* 8, 413–427.